



HELLENIC REPUBLIC

**National and Kapodistrian  
University of Athens**

EST. 1837

SCHOOL OF HEALTH SCIENCES  
DEPARTMENT OF DENTISTRY



Post-Graduate Program

"Community Dentistry, Prevention and Oral Health"

Specialization in Orthodontics

Title of Master's Thesis

**The effect of mandibular third molars on crowding of  
mandibular teeth, in patients with or without previous  
orthodontic treatment: a systematic review and meta-  
analysis.**

**Georgia Palikaraki**

Athens 2023

**Supervisor of Master's Thesis:**

Iosif Sifakakis, Associate professor



ΕΛΛΗΝΙΚΗ ΔΗΜΟΚΡΑΤΙΑ

**Εθνικόν και Καποδιστριακόν  
Πανεπιστήμιον Αθηνών**

———— ΙΔΡΥΘΕΝ ΤΟ 1837 ————

ΣΧΟΛΗ ΕΠΙΣΤΗΜΩΝ ΥΓΕΙΑΣ

ΤΜΗΜΑ ΟΔΟΝΤΙΑΤΡΙΚΗΣ



Πρόγραμμα Μεταπτυχιακών Σπουδών  
«Κοινωνική Οδοντιατρική, Πρόληψη και Στοματική Υγεία»

Ειδίκευση: ΟΡΘΟΔΟΝΙΚΗ

**Τίτλος Μεταπτυχιακής Διπλωματικής Εργασίας**  
**Ο ρόλος των τρίτων γομφίων της κάτω γνάθου στον πρόσθιο**  
**συνωστισμό της κάτω γνάθου, σε ασθενείς με/χωρίς**  
**προηγούμενη ορθοδοντική θεραπεία : συστηματική ανασκόπηση**  
**και μετά ανάλυση.**

**Γεωργία Παληκαράκη**

Αθήνα 2023.

**Επιβλέπων Καθηγητής  
για την εκπόνηση  
της Μεταπτυχιακής Διπλωματικής Εργασίας:**

Ιωσήφ Σηφακάκης, Αναπληρωτής Καθηγητής

## Ευχαριστίες

*Με αφορμή την ολοκλήρωση της μεταπτυχιακής μου εργασίας και την ολοκλήρωση του κύκλου σπουδών μου για την απόκτηση του μεταπτυχιακού τίτλου για την ειδικότητα της Ορθοδοντικής, θα ήθελα να ευχαριστήσω εγγράφως όλο το επιστημονικό επιτελείο του τμήματος.*

*Η ολοκλήρωση της μεταπτυχιακής μου εργασίας θα ήταν αδύνατη χωρίς την πολύτιμη υποστήριξη του καθηγητή μου, Αν. Καθηγητή του τμήματος Ορθοδοντική κ. Ιωσήφ Σηφακάκη. Του εκφράζω ένα βαθύ ευχαριστώ για όλη την υποστήριξη που μου προσέφερε τόσο σε κλινικό όσο και σε επιστημονικό επίπεδο.*

*Ιδιαίτερες ευχαριστίες θα ήθελα να εκφράσω στον Καθηγητή του τμήματος κ.Χαλαζωνίτη για την άριστη συνεργασία που είχαμε κατά την διάρκεια των μεταπτυχιακών μου σπουδών αλλά και στους υπόλοιπους καθηγητές κ.Βασταρδή, κ.Καναβάκη για την καθοδήγησή τους καθόλη την διάρκεια των σπουδών μου.*

*Επιπλέον οφείλω τις πιο ειλικρινείς μου ευχαριστίες στον Αν.Καθηγητή του τμήματός μας κ.Απόστολο Τσολάκη που πάντα με ενθάρρυνε σε επιστημονικό, κλινικό αλλά και προσωπικό επίπεδο.*

*Τέλος, θέλω να ευχαριστήσω του γονείς μου Νίκο και Μαρία και την αδερφή μου Ιωάννα για την αμέριστη συμπαράσταση και κατανόησή τους, οι οποίοι υπήρξαν πάντα ένα ανεκτίμητο στήριγμα για μένα και στους οποίους οφείλω όλη την διαδρομή των σπουδών μου μέχρι σήμερα αλλά και τον σύντροφό μου Δημήτρη για όλη την υπομονή, συμπαράσταση και υποστήριξή του.*

*Στους γονείς μου Νίκο και Μαρία.*

<b>Περίληψη.....</b>	<b>8</b>
<b>Abstract.....</b>	<b>10</b>
<b>Introduction.....</b>	<b>11</b>
Specific objectives or hypotheses.....	12
<b>Materials and Methods.....</b>	<b>12</b>
Protocol and Registration.....	12
Eligibility	12
Information Sources and Search Strategy.....	12
Study Selection.....	13
Risk of Bias in Individual Studies.....	13
Data Collection and Data Items.....	13
Summary Measures and Synthesis of the Results.....	14
Additional Analyses and Risk of Bias across Studies.....	14
<b>Results.....</b>	<b>15</b>
Study Selection.....	15
General Characteristics of the Included Studies.....	15
Quantitative Synthesis of Included Studies-Synthesis of Results.....	16
<b>Discussion.....</b>	<b>17</b>
Main findings in the context of the existing evidence and	17
Limitations.....	18
<b>Conclusions.....</b>	<b>19</b>
<b>References.....</b>	<b>20</b>
Tables.....	23
Figures.....	26

## ΠΕΡΙΛΗΨΗ

**Σκοπός:** Η συμμετοχή των κάτω τρίτων γομφίων στον δευτερογενή συνωστισμό και την κλίση των δοντιών παραμένει αμφιλεγόμενη. Έχει παρατηρηθεί αύξηση του συνωστισμού στους εφήβους και τους νέους ενήλικες μεταξύ 13-26 ετών, κατά την περίοδο ανατολής των τρίτων γομφίων. Πολλοί ορθοδοντικοί συνήθιζαν να παραπέμπουν τους ασθενείς τους να αφαιρέσουν προληπτικά τους τρίτους γομφίους προς αποφυγή του δευτερογενή συνωστισμού των πρόσθιων δοντιών και την υποτροπή της ορθοδοντικής θεραπείας. Ωστόσο έχει βρεθεί επίσης ότι ο δευτερογενής συνωστισμός είναι ένα πολυπαραγοντικά φαινόμενο και ότι οι αιτιολογικοί του παράγοντες διαφέρουν μεταξύ των ατόμων.

Ο σκοπός της παρούσας συστηματικής ανασκόπησης ήταν η αξιολόγηση της επιστημονικής τεκμηρίωσης σχετικά με τον ρόλο των τρίτων γομφίων της κάτω γνάθου στο συνωστισμό των πρόσθιων δοντιών, σε ασθενείς με ή/και χωρίς προηγούμενη ορθοδοντική θεραπεία. Οι δευτερεύουσες εκβάσεις (secondary outcomes) περιελάμβαναν μεταβολές στο εύρος και μήκος του κάτω οδοντικού τόξου.

**Υλικά και Μέθοδοι:** Η αναζήτηση διενεργήθηκε στις βάσεις δεδομένων PubMed, Scopus, Proquest και Google Scholar, μέχρι τον Απρίλιο 2022. Συμπεριελήφθησαν μελέτες που αξιολογούσαν τον ρόλο των τρίτων γομφίων της κάτω γνάθου στον συνωστισμό των κάτω πρόσθιων δοντιών, σε ασθενείς οποιασδήποτε ηλικίας και φύλου, με παρόντες τρίτους γομφίους της κάτω γνάθου (έγκλειστους, ημιέγκλειστους, που έχουν ανατείλει). Για την καταγραφή των απαραίτητων πληροφοριών χρησιμοποιήθηκαν προκαθορισμένα έντυπα συλλογής δεδομένων στα οποία καταγράφονταν οι εξής πληροφορίες: ο πρώτος συγγραφέας, η ηλικία, το είδος της μελέτης, αν είχε γίνει προηγούμενη ορθοδοντική θεραπεία, το μέγεθος του δείγματος, η παρουσία τρίτων γομφίων, τα εργαλεία ανάλυσης, ο τρόπος μέτρησης του συνωστισμού, τα αποτελέσματα της μελέτης. Η εκτίμηση του κινδύνου μεροληψίας έγινε με την χρήση του "ROBINS-I" (RoB) assessment tool .

**Αποτελέσματα:** Συνολικά 13 μελέτες συμπεριελήφθησαν στην παρούσα συστηματική ανασκόπηση. Οι περισσότερες από αυτές παρουσίαζαν μέτριο κίνδυνο μεροληψίας ενώ οι



υπόλοιπες υψηλό. Ο συνολικός αριθμός συμμετεχόντων ήταν 1342 από τις συμπεριλαμβανόμενες μελέτες. Όλες οι μελέτες που συμπεριλήφθηκαν ήταν μελέτες παρατήρησης. Για την μέτρηση του συνωστισμού χρησιμοποιήθηκαν πανοραμικές, πλάγιες κεφαλομετρικές ακτινογραφίες, εκμαγεία και αξονική τομογραφία κωνικής δέσμης. Ο συνωστισμός ποσοτικοποιήθηκε με τον δείκτη του Little στις περισσότερες μελέτες ενώ ορισμένες χρησιμοποίησαν την ανάλυση Lundstrom ή την αναλογία Ganss. Σε 4 μελέτες βρέθηκε συσχέτιση μεταξύ της παρουσίας κάτω τρίτων γομφίων και συνωστισμού των κάτω πρόσθιων δοντιών. Επτά από αυτές τις μελέτες συμπεριελήφθησαν στην ποσοτική ανάλυση. Πραγματοποιήθηκαν τρεις διαφορετικές μετα-αναλύσεις: για ασθενείς α) με ή β) χωρίς προηγούμενη ορθοδοντική θεραπεία και γ) σε συνδυασμό για ασθενείς με και χωρίς προηγούμενη ορθοδοντική θεραπεία. Σύμφωνα με τα συγκεντρωτικά αποτελέσματα και των τριών μετα-αναλύσεων, το μοντέλο τυχαίων επιδράσεων βρήκε στατιστικά σημαντικό όφελος για όσους δεν έχουν τρίτους γομφίους έναντι αυτών με τρίτους γομφίους, όσον αφορά στο συνωστισμό, στον δείκτη Little και στο μήκος του κάτω οδοντικού τόξου.

**Συμπεράσματα:** Οι κάτω τρίτοι γομφίοι πιθανόν να συμβάλλουν στη δημιουργία συνωστισμού της κάτω γνάθου και στη μείωση του μήκους του κάτω τόξου. Απαιτείται περαιτέρω έρευνα για να διευκρινιστεί ο ρόλος των τρίτων γομφίων στο συνωστισμό των προσθίων δοντιών της κάτω γνάθου.

*Η παρούσα διπλωματική εργασία έχει δημοσιευτεί στο επιστημονικό ορθοδοντικό περιοδικό *The Angle Orthodontist*.*

## **ABSTRACT**

**Objectives:** To map the scientific evidence related to the role of the mandibular third molars on the late crowding of the lower anterior teeth in patients with or/and without previous orthodontic treatment. The secondary outcomes included changes in mandibular arch width and arch length.

**Materials and Methods:** The databases PubMed, Scopus, Proquest and Google Scholar were searched for inception until April 2022. The included papers were studies evaluating the role of mandibular third molars on crowding of mandibular anterior teeth in patients of any age and gender, with mandibular third molars impacted or semi-impacted or erupted. Pre-determined and pre-piloted data collection forms were used to record the necessary information.

**Results:** 13 observational studies were included in the present systematic review. Most of them were assigned an overall risk of bias in terms of moderate risk while the rest of them were at high risk. Four studies found an association between the presence of mandibular third molar and mandibular incisor crowding. Finally, 7 studies were included in the quantitative analysis. Three different meta-analyses were conducted: for patients a) with or b) without previous orthodontic treatment and c) in combination for patients with and without previous orthodontic treatment. According to the pooled results of all three meta-analyses, random effects model yielded a significant benefit for those without third molars against those with third molars, regarding crowding, mean Little's irregularity index and mean arch length.

**Conclusions:** Lower third molars may contribute to mandibular crowding and lower arch constriction. Further prospective research of high quality is needed to clarify the impact of third molars on anterior mandibular crowding.

**KEY WORDS:** Anterior crowding; Tertiary crowding; Third molar; Wisdom teeth; Relapse; Systematic review

## INTRODUCTION

Dental crowding is a common orthodontic problem, defined as the discrepancy between tooth size and available space for proper alignment. It may be classified into primary, secondary and tertiary crowding.<sup>1</sup> Tertiary crowding can also be found in the literature as “late dental crowding”, since it appears mostly in the mandibular incisors during adolescence or later.<sup>2,3</sup> In patients who had received previous orthodontic treatment, relapse of primary crowding may also occur during this life period.<sup>4,5</sup>

Late crowding of the mandibular incisors has been reported in both treated and untreated individuals and is particularly common: half of the general population may present moderate to severe crowding.<sup>6,7</sup> This is a major concern among older orthodontic patients, since with increasing age, there is a gradual decrease in the exposure of the upper anterior teeth, and an increase in the exposure of the lower incisors. As a result, the crowding of the lower anterior teeth becomes more visible and has a greater impact on the aesthetics of the smile.<sup>8</sup> The mandibular dental arch becomes more square-shaped with increasing age and crowding tends to increase, due to a decrease in arch length and perimeter.<sup>9-13</sup>

Late mandibular crowding is a multifactorial phenomenon and its aetiological factors may differ among the individuals.<sup>9</sup> It is often attributed to a disturbance of the equilibrium between the dentition and the pressure of the tongue, lips, cheeks. In this case, the teeth tend to move until a new state of equilibrium is reached.<sup>10-11</sup> The effect of the mandibular third molars on late crowding remains controversial.<sup>3,4</sup> It has been reported that the lower third molars may generate a medial pressure that moves the lower posterior teeth forward, causing mandibular anterior crowding.<sup>14-16</sup> An increase in mandibular crowding has been reported in adolescents and young adults between the ages of 13-26 years, during the eruption of the third molars.<sup>12,17</sup> On the other hand, other reports claim that this pressure is not capable of causing anterior crowding.<sup>1,15,16</sup> Nowadays, some orthodontists refer their patients to an oral surgeon for third molar extraction in order to avoid re-crowding of the anterior teeth. A recent systematic review (SR) with meta-analysis evaluated only patients after previous orthodontic treatment and suggested mandibular third molar removal for preventing or alleviating long-term incisor irregularity.<sup>18</sup> Whether third molars contribute to anterior mandibular crowding, and the relation with orthodontic treatment remains controversial.

The present SR was conducted to answer the research question: “is there any association between the presence of lower third molars and late crowding of the lower anterior teeth?”

The objective was to map the scientific evidence related to the role of the mandibular third molars on the late crowding of the lower anterior teeth in patients with or/and without previous orthodontic treatment. Secondary outcomes included changes in mandibular arch width and arch length.

## **MATERIALS AND METHODS**

### **Protocol and Registration**

This SR was conducted according to a specific protocol developed and piloted in respect to the guidelines outlined in the Preferred Reporting Items for Systematic review and Meta-Analysis Protocols (PRISMA-P).<sup>19</sup> The study was registered at PROSPERO (CRD42023331648).

### **Eligibility Criteria**

Eligibility criteria were based on the PICOS (Participants, Intervention, Comparison, Outcomes, Study design) strategy (Table 1). Patients of any age and sex, in permanent dentition, with or without previous orthodontic treatment, presented with impacted / semi-impacted / erupted mandibular third molars were compared with patients with absence of mandibular third molars (agenesis / extracted).

Finite element analyses, follow-ups, case reports, reviews (systematic and literature), author's responses, letters to the editor, editor's summary, books and/or book chapters, abstracts, congress abstracts, personal opinions, authors debates, summary articles, non-English articles, or no author response to inquiry email for data clarification were excluded.

### **Information Sources and Search Strategy**

The search databases included PubMed, Scopus, ProQuest and Google Scholar. The aforementioned databases were searched from inception until April 30, 2022. The research was specified and performed to identify any relevant study based upon various combinations of key words (Table 2).

Developed detailed search strategies for each database were conducted and assessed with the help of a second investigator (Assist. Prof. Anastasia Mitsea). The search strategies developed for MEDLINE were adopted, nevertheless, they were revised appropriately for each database in order to include the differences in controlled vocabulary and syntax rules (Table 2). No restrictions were applied on the date and status of publication, except for the language: only English language papers were considered for inclusion. The reference lists of all the eligible studies, as well as these of relevant reviews, were searched. The authors of retrieved studies were contacted for further clarifications, if needed.

### **Study Selection**

Titles and abstracts of the retrieved records were assessed for inclusion independently and in duplicate with the help of a second investigator (Assist. Prof. Anastasia Mitsea). We were not blinded to the identity of the authors, their institution, or the results of the research. Any article title that met the objectives of the study but did not have an available abstract was fully analyzed in the final evaluation. Full texts of the selected studies were obtained and evaluated to verify whether they fulfilled the eligibility criteria. Disagreements were resolved by discussion or consultation with a third investigator before the final decision (Assoc. Prof. Iosif Sifakakis). The reference lists of all retrieved full text articles were searched for relevant articles that might have been missed through the electronic search. Finally, the articles that didn't answer the clinical research questions were excluded.

### **Risk of Bias in Individual Studies**

We assessed the risk of bias in individual studies, independently and in duplicate, using the "ROBINS-I" (RoB) assessment tool. Any disagreements were resolved by discussion or consultation with the third investigator (Assoc. Prof. Iosif Sifakakis).<sup>20</sup>

### **Data Collection and Data Items**

Data extraction was performed independently and in duplicate with the help of a second investigator (Assist. Prof. Anastasia Mitsea). Any disagreements were resolved through discussion with the third investigator (Associate Professor Iosif Sifakakis). Predetermined and pre-piloted data collection forms were used to record the necessary information: first author;

age; type of study; previous orthodontic treatment; sample size; presence of third molars; analysis tools; crowding measurement; angulation of lower teeth; outcome of the study.

### **Summary Measures and Synthesis of the Results**

The summary characteristics of the included studies were reviewed. The mean values and the corresponding SDs were extracted independently from each article that used quantitative measures for outcome assessment. The random effects model was used. Forest plots were also used to visualize the pooled estimates across studies. Statistical analysis of the data was conducted with the statistical package R ([www.r-project.org](http://www.r-project.org), v3.6.2) while the level of statistical significance was set at 0.05.

### **Additional Analyses and Risk of Bias across Studies**

Egger's test was used to assess potential publication bias in the meta-analysis via funnel plot asymmetry. If there were at least two studies with the same outcome and having sufficient homogeneity regarding the compared groups and the time frame, meta-analysis would be undertaken. Otherwise, a qualitative synthesis of the results would be reported. Statistical heterogeneity was inspected with the DerSimonian and Laird's Q statistic and I<sup>2</sup> statistic. If split-mouth studies were included in meta-analysis, the correlation coefficient was planned to be computed, in order to calculate the adjusted standard error (Se), as described in Chapter 23 in Cochrane handbook. The difference in means of both arms and the adjusted standard error were used in the meta-analysis.<sup>21</sup>

## RESULTS

### Study Selection

The results of literature searching identification, inclusion and exclusion of the articles are presented in the flow diagram according to the PRISMA statement (Figure 1). Initially, 847 relevant records were identified after the electronic and manual search, while only 798 remained after manual duplicate removal. After the title and abstract screening, 27 articles were selected for full-text review, according to the inclusion and exclusion criteria. Finally, 9 papers were excluded because 4 of them were reviews, 2 were case series, 1 pilot study, 1 questionnaire and 1 unclassified. The remaining 5 did not meet the inclusion criteria for outcome and comparator. Finally, 13 articles were included in the qualitative synthesis.<sup>1,22-33</sup> However, only 7 articles were included in the quantitative synthesis since the remaining studies were heterogeneous.<sup>1,22-24,26,28,30</sup>

### Risk of Bias in Individual Studies

Risk of bias for the included studies<sup>1,22-33</sup> is depicted in Supplemental Table 1 and Table 3 (in summary). 8 studies were assigned an overall risk of bias in terms of moderate risk<sup>1,23,25,29-31,33</sup> while the rest of them (5 studies) in terms of high risk.<sup>22,26-28,32</sup>

### General Characteristics of the Included Studies

The general characteristics of the studies included in the present SR, as well as the sample characteristics are depicted in Tables 4 and 5. A total number of 1342 participants was calculated from the included studies. All the included studies were observational by design (cross sectional and prospective non-randomized controlled trials)<sup>1,22,23-33</sup> and used 2D radiographs (orthopantomographs, lateral cephalometric radiographs) and casts as analysis tools, except two studies, that used only casts<sup>29</sup> or cone beam computed tomography (CBCT)<sup>30</sup>. Crowding was quantified with Little's irregularity index in 8 studies<sup>23,22,26-30,33</sup>, 3 used Lundstrom analysis<sup>1,20,31</sup> and 2 studies used Ganss Ratio.<sup>31,33</sup> Five studies measured arch length<sup>22,26-28,32</sup> and 4 studies measured arch width<sup>1,25,26,28</sup>. In 4 studies the participants had undergone orthodontic treatment.<sup>22,26-28</sup> Four out of the 13 eligible studies supported a cause-and-effect relationship among third molars and anterior crowding.<sup>29-32</sup>

### **Quantitative Synthesis of Included Studies-Synthesis of Results**

Seven studies were included in the quantitative analysis and three different meta-analyses were conducted. The first one was conducted in papers without previous orthodontic treatment,<sup>1,23,24,30</sup> and the second in patients with previous orthodontic treatment.<sup>22,26,28</sup> The third analysis was implemented in combination for patients with and without previous orthodontic treatment.<sup>1,22-24,26,28,30</sup>

The first meta-analysis of 4 studies was performed on patients without previous orthodontic treatment and compared mean differences in crowding between cases with or without third molars (agenesis or extracted). Irregularity was statistically significant with random-effects model (\*\* $P=.01$ ). According to the pooled results of the meta-analysis, the random effects model yielded a significant benefit for those without third molars against those with third molars (Figure 2). Specifically, those with third molars had greater mean Little's irregularity index scores in comparison with those without third molars. Egger's test  $p$ -value was 0.09 which implies borderline significant publication bias.

The second meta-analysis of 3 studies on patients with previous orthodontic treatment, compared mean differences in cases with and without third molars (agenesis or extracted), regarding crowding, arch width and arch length. The pooled estimate of crowding and arch width was not statistically significant ( $P=.23$ ,  $P=.69$ ). (Figure 3). Moreover, Egger's test  $P$ -value was 0.73 which implies no publication bias.

The third meta-analysis of 7 studies on patients with and without previous orthodontic treatment, compared mean differences in crowding between patients with and without third molars (agenesis or extracted). Crowding was statistically significant with random effects model (\*\* $P=.005$ ). According to the pooled results of meta-analysis, random effects model yielded a significant benefit for those without third molars against those with third molars. Specifically, those with third molars had greater mean Little's irregularity index scores in comparison with those without third molars (Figure 4). Moreover, Egger's test  $P$ -value was 0.10 which implies no evidence of publication bias.



## DISCUSSION

Three SRs have been conducted to assess the possible association between mandibular third molars and lower anterior crowding,<sup>18,34,35</sup> however only the most recent one (2018) included a meta-analysis<sup>18</sup>. The first available SR assessed the literature by using the Medline database between 1971 and 2011 and included 21 articles in the qualitative analysis.<sup>35</sup> The results were contradictory: four researchers concluded that lower third molars favor teeth crowding, however this finding was not confirmed by the rest of the articles. The second was a systematic literature review that selected 12 observational studies to evaluate the role of third molars in late mandibular anterior crowding in post orthodontic patients. The qualitative analysis of the results failed to support the cause-and-effect relationship between third molars and lower anterior tooth crowding. However, according to these authors, definitive conclusions on the role of the third molars in the development of anterior tooth crowding cannot be drawn.<sup>34</sup> Finally, the third SR evaluated quantitatively with a meta-analysis three retrospective studies conducted on patients in the retention phase following orthodontic treatment. They found statistically significant differences between the erupted third molar extraction group and agenesis third molar group regarding the Little's irregularity index, whereas the arch length and intermolar width did not differ between these patient groups. These authors suggest mandibular third molar removal for preventing or alleviating long-term incisor irregularity.<sup>18</sup> The present SR, was the first to investigate the role of the mandibular third molars on the space conditions of the lower teeth in patients with or/and without previous orthodontic treatment.

The present SR identified no randomized trials on the effect of third molars on anterior teeth. The eligible studies were intervention studies with control group, without randomization / double blinding and most of them did not find such an association, suggesting a coincidence between the two events. These studies examined both orthodontically treated and untreated subjects, with either impacted, erupted, extracted, or congenitally absent (agenesis) third molars.<sup>1,22-28,33</sup> Four studies report a different outcome and support a cause-and-effect relationship among third molars and anterior crowding.<sup>29-32</sup>

These contradictory results may be explained by differences in study design and methodology, sample size, inclination and degree of impaction (impacted, semi impacted, erupted) of the third molars. The present meta-analyses suggest that third molars may contribute to mandibular crowding. However, the aetiology of this phenomenon is

multifactorial and the extraction of third molars per se may not prevent tertiary crowding or dental relapse after orthodontic treatment. Thus, removing third molars, whether impacted or not, only for preventing crowding is not justified or recommended by the present results. The extraction of third molars is indicated in cases of posterior crowding, if distalization is needed, or when their unfavorable orientation compromises the stability of the arch or the prognosis of the adjacent second molar.<sup>26,36</sup> Nevertheless, strict retention protocols should be utilized after orthodontic treatment in patients with lower third molars left in situ.

The extracted data were classified according to the presence or absence (agenesis/extracted) of the third molar and three meta-analyses were conducted: for patients with or without orthodontic treatment as well as in combination for patients with and without previous orthodontic treatment. According to the pooled results of the meta-analysis, random effects models yielded a significant benefit for those without third molars against those with third molars in the two meta-analyses. Specifically, patients with third molars had greater mean Little's irregularity index scores in comparison with those without third molars in the non-orthodontic patient group. When all patients (with and without previous orthodontic treatment) were pooled together, the meta-analysis revealed a statistically significant effect of the third molar on random effects model (\* $P=.02$ ).

### **Strengths and Limitations**

A protocol was developed a priori and the present review followed clear-cut guidelines. The search strategies employed were exhaustive, covering electronic, manual, and gray literature material up to April 2022. Their character was comprehensive, irrespective of date and status of publication. Every effort to decrease bias in the methodology employed was made. Screening, verification of eligibility, abstraction of information, assessment of risk of bias and of the quality of evidence were all performed in duplicate, and any disagreement was resolved by discussion or consultation until a final consensus was achieved.

However, a potential source of bias in the present review could be the exclusion of articles not written in English. Further limitations in this study arise mainly from the nature and the characteristics of the retrieved data during the review process. Factors such as extractions, interproximal enamel reduction or compliance with fixed/removable retention protocols were not evaluated in the included studies. However, the main limitation was related to the heterogeneity of the studies and the inadequate control of confounding in the included

studies. These were non randomized trials, with risk of bias and differences in the methods of assessing the crowding. Differences in the study groups (impacted / erupted / present third molars vs. agenesis / extracted) may have confounded the present findings and the impact of the third molars on the mandibular incisors may have occurred before their extraction. Moreover, each of the first and second meta-analyses included four studies. Egger's test may lack the statistical power to detect bias when the number of studies is small (<10). Additionally, there are several other factors that may affect anterior crowding, such as growth-related changes, muscular factors, periodontal ligament traction, bone adaptation, masticatory force etc.<sup>17,18</sup> Further research is necessary to clarify the etiology of mandibular anterior crowding and to quantify the relapse tendency in each individual case.

## **CONCLUSIONS**

- Within the limitations of the present systematic review, it can be concluded that lower third molars may contribute to mandibular crowding
- More specifically, the degree of lower anterior dental crowding was higher in patients with presence of mandibular third molars, at least in the included studies.

## REFERENCES

1. Sidlauskas A, Trakiniene G. Effect of the lower third molars on the lower dental arch crowding. *Stomatologija*. 2006;8:80-84.
2. van der Linden FP. Theoretical and practical aspects of crowding in the human dentition. *J Am Dent Assoc*. 1974;89:139-153.
3. Buschang PH, Shulman JD. Incisor crowding in untreated persons 15-50 years of age: United States, 1988-1994. *Angle Orthod*. 2003;73:502–508.
4. Hussain SS, Ashraf B, Khan S. Relationship of dental crowding to tooth size and arch dimensions in class I normal & class I malocclusion sample. *Pak Oral Dental J*. 2014;34:660-664.
5. Littlewood SJ, Millett DT, Doubleday B, Bearn DR, Worthington HV. Retention procedures for stabilising tooth position after treatment with orthodontic braces. *Cochrane Database Syst Rev*. 2016;29:CD002283.
6. Buschang PH, Stroud J, Alexander RG. Differences in dental arch morphology among adult females with untreated Class I and Class II malocclusion. *Eur J Orthod*. 1994;16:47–52.
7. Marques LS, Pordeus IA, Ramos-Jorge ML, Filognio CA, Filognio CB, Pereira LJ, Paiva SM. Factors associated with the desire for orthodontic treatment among Brazilian adolescents and their parents. *BMC Oral Health* 2009;9:34.
8. Dong JK, Jin T.H, Cho HW, Oh SC. The esthetics of the smile: A review of some recent studies. *Int J Prosthodont*. 1999;12:9–19.
9. Bishara SE. Third molars: a dilemma! Or is it? *Am J Orthod Dentofacial Orthop*. 1999;115:628–633.
10. Little RM, Wallen TR, Riedel RA. Stability and relapse of mandibular anterior alignment – first premolar extraction cases treated by traditional edgewise orthodontics. *Am J Orthod*. 1981;80:349-365.
11. Consolaro A, Cardoso MA. Mandibular anterior crowding: normal or pathological? *Dental Press J Orthod*. 2018;23:30-36.
12. Galic I, Vodanović M, Cameriere R, Nakaš E, Galić E, Selimović E, Brkić H. Accuracy of Cameriere, Haavikko, and Willems radiographic methods on age estimation on Bosnian-Herzegovian children age groups 6–13. *Int J Leg Med*. 2011;125:315–321.

13. Mazzilli LEN, Melani RFH, Lascala CA, Palacio LAV, Cameriere R. Age estimation: Cameriere's open apices methodology accuracy on a southeast Brazilian sample. *J Forensic Leg Med.* 2018;58:164–168.
14. Vego L. A longitudinal study of mandibular arch perimeter. *Angle Orthod.* 1962;32:187-192.
15. Richardson ME. The role of the third molar in the cause of late lower arch crowding: a review. *Am J Dentofacial Orthop.* 1989;95:79-83.
16. Almpani K, Kolokitha OE. Role of third molars in orthodontics. *World J Clin Cases.* 2015;3:132-140.
17. Al Qassar SS, Mavragani M, Psarras V, Halazonetis DJ. The anterior component of occlusal force revisited: direct measurement and theoretical considerations. *Eur J Orthod.* 2016;38:190-196.
18. Cheng HC, Peng BY, Hsieh HY, Tam KW. Impact of third molars on mandibular relapse in post-orthodontic patients: A meta-analysis. *J Dent Sci.* 2018;13:1-7.
19. Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA, PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ.* 2015;350:g7647.
20. Sterne JA et al. ROBINS-I: a tool for assessing risk of bias in a non-randomised studies of interventions. *BMJ.* 2016;355:i4919.
21. Higgins JPT, Green S. Cochrane handbook for systematic reviews of interventions, version 5.1.0.
22. Ades AG, Joondeph DR, Little RM, Chapko MK. A long-term study of the relationship of third molars to changes in the mandibular dental arch. *Am J Orthod Dentofac Orthop.* 1990;97:323–335.
23. Zigante M, Pavlic A, Morelato L, Vandevska-RadunovicV, Spalj S. Presence and maturation dynamics of mandibular third molars and their influence on late mandibular incisor crowding: a longitudinal study. *Int J Environ Res Public Health* 2021;18:10070.
24. Shah R, Kanzariya N, Goje S, Kulkarni N, Joshi H, Chellani S. Assessment of role of mandibular third molar position in lower anterior crowding- a cross sectional study. *J Integr Health Sci.* 2018;6:69–73.
25. Stanaityte R, Trakiniene G, Gervickas A. Lower dental arch changes after bilateral third molar removal. *Stomatologija.* 2014;16:31-36.

26. Harradine NW, Pearson MH, Toth B. The effect of extraction of third molars on late lower incisor crowding: a randomized controlled trial. *Br J Orthod.* 1998;25:117–22.
27. van der Schoot EA, Kuitert RB, van Ginkel FC, Prah-Andersen B. Clinical relevance of third permanent molars in relation to crowding after orthodontic treatment. *J Dent.* 1997;25:167–169.
28. Kaplan R.G. Mandibular third molars and postretention crowding. *Am J Orthod.* 1974;66:411–430.
29. Esan T, Schepartz LA. Third molar impaction and agenesis: influence on anterior crowding. *Ann Hum Biol.* 2017;44:46-52.
30. Husain S, Rengalakshmi S. Correlation between mandibular third molar and mandibular incisor crowding: a retrospective CBCT-based study. *J Dent Res Dent Clin Dent Prospects.* 2021;15: 247-250.
31. Niedzielska I. Third molar influence on dental arch crowding. *Eur J Orthod.* 2005;27:518–523.
32. Lindqvist B, Thilander B. Extraction of third molars in cases of anticipated crowding in the lower jaw. *Am J Orthod.* 1982;81:130–139.
33. Hasegawa Y, Terada K, Kageyama I, Tsuchimochi T, Ishikawa F, Nakahara S. Influence of third molar space on angulation and dental arch crowding. *Odontology.* 2013;101:22–28.
34. Zawawi KH, Melis M. The role of mandibular third molars on lower anterior teeth crowding and relapse after orthodontic treatment: a systematic review. *Sci World J.* 2014;615429.
35. Stanaityte R, Trakiniene G, Gervickas A. Do wisdom teeth induce lower anterior teeth crowding? A systematic literature review. *Stomatologia.* 2014;16:15e8.
36. Salagnac JM. L'indication de germectomie ou d'extraction des 3èmes molaires chez les sujets en cours ou en fin de traitement d'orthopédie dento faciale est-elle toujours justifiée? *Rev Orthop Dento Faciale.* 2014;48:249-260.

**Table 1.** Eligibility Criteria Based on the PICOS (Participants, Intervention, Comparison, Outcomes, Study Design) Strategy

Participants (population)	Patients of any age and sex in permanent dentition with or without previous orthodontic treatment
Intervention	Patients presented with mandibular third molars impacted or semi-impacted or erupted
Comparisons	Patients with absence of mandibular third molars (agenesis; extracted)
Outcome	(a) Primary: Little's irregularity index, Ganss ratio, Lundström analysis (b) Secondary: arch length, arch width Outcome measured: on casts, panoramic radiographs, lateral cephalometric radiographs, or cone beam computed tomography
Study design	Randomized clinical trials, non- or quasi-randomized controlled trials, prospective or retrospective studies evaluating the association of third molars and late anterior crowding or relapse

**Table 2.** Electronic Search Strategy

Databases	Search Strategy/Keywords
Medline/PubMed	(secondary crowding OR relapse OR anterior crowding OR incisor crowding OR mandibular crowding OR incisor relapse OR anterior post retention OR anterior post-retention OR re-crowding) AND (third molar OR third molars OR wisdom tooth OR wisdom teeth OR 3rd molar OR teeth wisdom OR tooth wisdom OR molar third)
Scopus	KEY ((secondary crowding OR relapse OR anterior crowding OR incisor crowding OR mandibular crowding OR incisor relapse OR anterior post retention OR anterior post-retention OR re-crowding) AND (third molar OR third molars OR wisdom tooth OR wisdom teeth OR 3rd molar))
Google Scholar	allintitle: (secondary crowding OR relapse OR anterior crowding OR incisor crowding OR mandibular crowding OR incisor relapse OR anterior post retention OR anterior post-retention OR re-crowding) AND (third molar OR third molars OR wisdom tooth OR wisdom teeth OR 3rd molar)
ProQuest Dissertations and Theses Global	Ti (secondary crowding OR relapse OR anterior crowding OR incisor crowding OR mandibular crowding OR incisor relapse OR anterior post retention OR anterior post-retention OR re-crowding) AND (third molar OR third molars OR wisdom tooth OR wisdom teeth OR 3rd molar)

**Table 3.** Risk of Bias Summary

Studies	Zigante et al., 2021 <sup>23</sup>	Husain et al., 2021 <sup>30</sup>	Shah et al., 2018 <sup>24</sup>	Esan et al., 2017 <sup>29</sup>	Stanaitytė et al., 2014 <sup>25</sup>
<b>Bias due to confounding</b>	Serious	Serious	Serious	Serious	Serious
<b>Bias in selection of participants into the study</b>	Low	Low	Low	Low	Low
<b>Bias in classification of interventions</b>	Low	Low	Low	Low	Low
<b>Bias due to departures from intended interventions</b>	Low	Low	Low	Low	Low
<b>Bias due to missing data</b>	Low	Low	Low	Low	Low
<b>Bias in measurement of outcomes</b>	Low	Low	Low	Low	Low
<b>Bias in selection of the reported result</b>	Low	Low	Low	Low	Low
<b>Overall RoB</b>	<b>Moderate</b>	<b>Moderate</b>	<b>Moderate</b>	<b>Moderate</b>	<b>Moderate</b>

<sup>a</sup> RoB indicates ROBINS-I assessment tool.

**Table 3.** Extended

Hasegawa et al., 2013 <sup>33</sup>	Sidlauskas & Trakiniene, 2006 <sup>1</sup>	Niedzielska, 2005 <sup>31</sup>	Lindqvist & Thilander, 1982 <sup>32</sup>	Harradine et al., 1998 <sup>26</sup>	Van der Scoot et al., 1997 <sup>27</sup>	Ades et al., 1990 <sup>22</sup>	Kaplan et al., 1974 <sup>28</sup>
Serious	Serious	Serious	Serious	Serious	Serious	Serious	Serious
Moderate	Low	Low	Moderate	Moderate	Moderate	Moderate	Moderate
Low	Low	Low	Low	Low	Low	Low	Low
Low	Low	Low	Low	Low	Low	Low	Low
Low	Low	Low	Low	Low	Low	Low	Low
Low	Low	Low	Low	Low	Low	Low	Low
Low	Low	Low	Low	Low	Low	Low	Low
<b>Serious</b>	<b>Moderate</b>	<b>Moderate</b>	<b>Serious</b>	<b>Serious</b>	<b>Serious</b>	<b>Serious</b>	<b>Serious</b>

Table 4. Summary Characteristics of Included Studies

		Type of Study	Age (y)	Sample Size (N)	Inclusion / Exclusion
Without previous orthodontic treatment	Zigante et al., 2021 <sup>23</sup>	Cross-sectional	12, 15, 18, and 21	72 (F: 34, M: 38)	No significant malocclusions nor facial disharmonies at the start of follow-up (6 y), no orthodontic treatment. Presence of panoramic radiographs and plaster casts at the ages of 12, 15, 18, and 21
	Husain et al., 2021 <sup>30</sup>	Cross-sectional	18–30	40 patients in 2 groups: (a) without third molars (20) and (b) with third molars (20)	Inclusion criteria: Class I malocclusion, CBCT images of patients in the 18–30 age group, CBCT images recorded in the database of the University, CBCT images with the presence of all the permanent dentition Exclusion criteria: presence of impacted teeth, presence of deciduous teeth, presence of skeletal asymmetries, presence of any underlying pathologies, partial or complete absence of teeth, other than third molars
	Shah et al., 2018 <sup>24</sup>	Cross-sectional	>25	90 (F 50, M 40) in 3 groups: (a) group E with erupted M3: 58, (b) group UE with unerupted M3: 18, and (c) group AB with agenesis of M3: 14	Exclusion criteria: patients with a history of previous orthodontic, orthopedic, or facial and surgical treatment; systemic disease, developmental anomalies, or syndromes; abnormal habits and third molar extraction
	Esan et al., 2017 <sup>29</sup>	Cross-sectional	M 19–70 (43.66 SD 12.00)	535 in 6 groups: (a) bilateral M3 present: 439, (b) unilateral agenesis: 23, (c) bilateral agenesis: 14, (d) unilateral impaction: 36, (e) bilateral impaction: 21 and (f) agenesis and impaction: 2	Inclusion criteria: Black South African males
	Stanaitytė et al., 2014 <sup>25</sup>	Prospective non randomized controlled trial	25,5 (16,2–55,1)	30 (F 19, M 11) before and after lower M3 removal	<ul style="list-style-type: none"> <li>• complete lower dental arch</li> <li>• age at least 16 y</li> <li>• no orthodontic treatment before records collected</li> <li>• bilateral LM3 removal</li> <li>• good quality OPT and plaster casts</li> </ul>
	Hasegawa et al., 2013 <sup>33</sup>	Cross-sectional	21,0 (18,3–24,1)	34 in 2 groups: (a) Class I normal occlusion (<3.5 mm) with F 10 and M 4, and (b) Class I crowding (≥3.5 mm) with F 10 and M 10)	Inclusion criteria: Angle Class I molar relationship, all four M3 impacted, all teeth were caries-free, no previous dental treatment, no anomalies of crown morphology, no orthodontic treatment in either maxillary or mandibular arch
	Sidlauskas & Trakinienė, 2006 <sup>1</sup>	Cross-sectional	21,01 (SD 4,13)	91	Inclusion criteria: age at least 17 y, complete lower dental arch (except M3), no orthodontic treatment before records collected, good state of care of the lower teeth with no artificial dental crowns, good quality OPTs and plaster casts available
Previous orthodontic treatment	Niedzielska, 2005 <sup>31</sup>	Cross-sectional	14–32	47 (F 36, M 11) in 4 groups: (a) bilaterally extracted: 17, (b) unilaterally extracted: 12, (c) control group (bilaterally retained): 16 and (d) M3 agenesis: 2	Exclusion criteria: previous orthodontic treatment or patients presented with malocclusions
	Lindqvist & Thilander, 1982 <sup>32</sup>	Cross-sectional	15,5 (13–19), with a 3 y follow-up	52	
	Harradine et al., 1998 <sup>26</sup>	Cross-sectional	14,8 (SD 16,2)	77 (F 45, M 32) in 2 groups: (a) M3 extracted: 44 and (b) M3 nonextracted: 33	Inclusion criteria: Class I malocclusion, CBCT images of patients in the 18–30 y age group, CBCT images recorded in the database of the University, CBCT images with the presence of all permanent dentition. Exclusion criteria: presence of impacted or deciduous teeth, presence of skeletal asymmetries, presence of any underlying pathologies, partial or complete absence of teeth, other than third molars
	Van der Scoot et al., 1997 <sup>27</sup>	Cross-sectional	22,3 (SD 4,2)	99 (F 60, M 39) in 4 groups: (a) both M3 erupted (lower arch n = 24, upper arch n = 23), (b) neither of M3 erupted (lower arch n = 19, upper arch n = 22), (c) both M3 extracted (lower arch n = 43; upper arch n = 37 and (d) one or both M3 congenitally absent (lower arch n = 8, upper arch n = 7)	-
	Ades et al., 1990 <sup>22</sup>	Cross-sectional	28,6 (18,6–39,4)	97 in 4 groups: (a) erupted: 32, (b) impacted: 14, (c) extracted: 34 and (d) absent: 17	1. All participants were Caucasian 2. Participants free of all retention for at least 10 y
Kaplan et al., 1974 <sup>28</sup>	Cross-sectional	26.6 (at postretention)	75 in 3 groups: (a) erupted: 30, (b) impacted: 20 and (c) agenesis: 25	Orthodontically treated Caucasian patients	

<sup>a</sup> CBCT indicates cone beam computed tomography and OPT orthopantomogram.



**Table 4. Extended**


Groups	Analysis Tools	Outcomes	Results
(a) agenesis (b) extracted (c) nonerupted (d) impacted	OPT casts	Little's irregularity index	No difference
(a) without third molars (b) with third molars	CBCT	Little's irregularity index	Association between the presence of mandibular third molar and mandibular incisor crowding
(a) erupted (visible in oral cavity, either partially or completely), (b) unerupted (not visible in oral cavity), (c) agenesis (neither visible in oral cavity nor in OPT)	OPT casts	Lundström analysis	No difference
(a) bilateral M3 present (b) unilateral agenesis (c) bilateral agenesis (d) unilateral impaction (e) bilateral impaction and (f) agenesis and impaction	Casts (jaws)	Little's irregularity index	Third molar impaction plays a role in anterior crowding
(a) Before M3 removal and (b) after M3 removal	OPT casts	Lower dental arch width changes	No difference
(a) without and (b) with crowding	OPT casts Cephalogram	Little's irregularity index , Ganss ratio	No difference
(a) erupted M3, (b) unerupted M3 and (c) M3 agenesis	OPT casts	Mesiodistal width related to the length from central incisor to canine, Lundström analysis	No difference
(a) bilaterally extracted (b) unilaterally extracted (c) bilaterally retained (d) M3 agenesis	OPT casts	Lundström's method, Ganss ration	Retained M3 associated with increased tooth crowding in relation to Ganss ratio
	Casts Cephalograms	Arch length, Cephalometric analysis	Extraction side had a more favorable development than the control side on 70% of the patients
(a) M3 extracted (b) M3 non-extracted	OPT casts Cephalograms	Little's irregularity index, intercanine width and arch length; cephalometric analysis	No difference
(a) right and left M3 had emerged, (b) neither of M3 had erupted, (c) right and left M3 were extracted, (d) one or both M3s were congenitally absent	OPT casts	Little's irregularity index, Arch length	No difference
(a) erupted, (b) impacted, (c) extracted, (d) agenesis	Casts Cephalograms	Little's irregularity index Arch length	No difference
(a) bilaterally erupted, in occlusion (b) bilaterally impacted (c) bilateral agenesis	Casts Cephalograms	Little's irregularity index Arch length Intermolar width	No difference

**Table 5.** General Characteristics of the Studies Included in the Quantitative Synthesis

	Groups	Number		
		With M3	Without M3	
<b>Without previous orthodontic treatment</b>	<b>Zigante et al., 2021<sup>23</sup></b>	(a) agenesis, (b) extracted, (c) impacted, (d) unerupted	present 48 (impacted at 21 y: 16; unerupted at 21 y: 10; erupted at 21 y: 22)	extracted: 16 hypodontia: 8
	<b>Husain et al., 2021<sup>30</sup></b>	(a) without M3 (b) with M3	20	20
	<b>Shah et al., 2018<sup>24</sup></b>	E: erupted (visible in oral cavity, either partially or completely) UN: unerupted (not visible in oral cavity) AB: agenesis (neither visible in oral cavity nor in OPT)	E:119 UE:38	AB: 23
	<b>Sidlauskas &amp; Trakiniene, 2006<sup>1</sup></b>	E: erupted M3 UE: unerupted M3 A: agenesis of M3	E: 88 UE: 85	A: 9
<b>Previous orthodontic treatment</b>	<b>Harradine et al., 1998<sup>26</sup></b>	3EX: extracted M3 3NEX: non-extracted M3	3NEX: 33	3EX: 44
	<b>Ades et al., 1990<sup>22</sup></b>	Impacted Erupted Extracted Agenesis	Erupted: 32 Impacted: 14	Absent: 17 Extraction: 34
	<b>Kaplan et al., 1973<sup>28</sup></b>	Bilaterally erupted: M3E Bilaterally impacted: M3I Bilateral agenesis: M3A	Erupted: 30 Impacted: 20	Agenesis 25

**Table 5.** Extended

Crowding (mm)		Arch Width (mm)		Arch Length (mm)	
With M3 Mean (SD)	Without M3 Mean (SD)	With M3	Without M3	With M3	Without M3
present: 1,8 (1,6) (impacted at 21 y: 1,3 (1,4), unerupted at 21 y: 1,9 (1,5), erupted at 21 y 1,7 (1,6)) group b: 6,79 (5,45)	extracted: 1,7 (2,0), agenesis: 0,5 (1,8) group a: 4,26 (4,88)	-	-	-	-
right side: E 2,02 (1,42), UE 1,3 (0,68), left side: E 1,6 (1,46), UE 1,4 (0,89)	right side: AB 1,5 (0,66), left side: 1,32 (0,57)	-	-	-	-
E: 1,50 (1,77) UE: 1,70 (2,20)	A: 0,78 (1,13)	-	-	-	-
3NEX: 1,10 (2,72)	3EX: 0,80 (1,23)	3NEX: -0,38 (0,38)	3EX: 0,37 (0,73)	3NEX: -2,13 (0,96)	3EX: -1,10 (1,13)
Impacted: 2,27 (1,81) Erupted: 3,19 (2,20)	Extracted: 3,25 (5,24) Agenesis: 2,55 (1,40)	Impacted: -1,52 (1,23) Erupted: -1,71 n(1,45)	Extracted: -1,86 (1,29) Agenesis: -1,47 (1,76)	Impacted: -2,01 (1,41) Erupted: -2,81 (2,12)	Extracted: -2,02 (1,71) Agenesis: -2,17 (2,10)
M3E: 3,00 (2,15) M3I: 3,35 (2,20)	M3A: 1,99 (1,76)	M3E: -1,38 (1,19) M3I: -1,49 (1,24)	M3A: -1,94 (1,27)	M3E: -2,50 (1,57) M3I: -2,39 (1,83)	M3A: -2,20 (1,55)



**Figure 1.** Flow chart diagram.




Figure 2. Forest plot for studies on patients without previous orthodontic treatment for crowding.




Figure 3. Forest-plot for studies on patients with previous orthodontic treatment for arch length.




Figure 4. Forest plot for all the included studies (on patients with and without previous orthodontic treatment) for crowding.



